

PROOF OF A CONJECTURE OF KLEINBERG-SAWIN-SPEYER

LUKE PEBODY

ABSTRACT. In Ellenberg and Gijswijt's groundbreaking work [2], the authors show that a subset of $\mathbb{Z}_3^n$ with no arithmetic progression of length 3 must be of size at most 2.755^n (no prior upper bound was known of $(3 - \epsilon)^n$), and provide for any prime p a value $\lambda_p < p$ such that any subset of $\mathbb{Z}_p^n$ with no arithmetic progression of length 3 must be of size at most λ_p^n .

Blasiak et al [1] showed that the same bounds apply to tri-coloured sum-free sets, which are triples $\{(a_i, b_i, c_i) : a_i, b_i, c_i \in \mathbb{Z}_p^n\}$ with $a_i + b_j + c_k = 0$ if and only if $i = j = k$.

Building on this work, Kleinberg, Sawin and Speyer [3] gave a description of a value μ_p such that no tri-coloured sum-free sets of size $e^{\mu_p n}$ exist in $\mathbb{Z}_p^n$, but for any $\epsilon > 0$, such sets of size $e^{(\mu_p - \epsilon)n}$ exist for all sufficiently large n . The value of μ_p was left open, but a conjecture was stated which would imply that $e^{\mu_p} = \lambda_p$, i.e. the Ellenberg-Gijswijt bound is correct for the sum-free set problem.

The purpose of this note is to close that gap. The conjecture of Kleinberg, Sawin and Speyer is true, and the Ellenberg-Gijswijt bound is the correct exponent for the sum-free set problem.

1. INTRODUCTION AND DEFINITIONS

Let G be an abelian group. Define a *tri-coloured sum-free set* in G^n to be a collection of triples (a_i, b_i, c_i) in G^n such that $a_i + b_j + c_k = 0$ if and only if $i = j = k$. Write $\text{sfs}(G^n)$ for the size of the largest sum-free set in G^n and $r_3(G^n)$ for the size of the largest set in G^n with no three-term arithmetic progression. If $X \in G^n$ is a set with no three-term arithmetic progression, then $\{(x, x, -2x) : x \in X\}$ is a sum-free set, so $r_3(G^n) \leq \text{sfs}(G^n)$.

Kleinberg, Sawin and Speyer [3] found a tight bound for $\lim \text{sfs}(\mathbb{Z}_p^n)^{\frac{1}{n}}$. To state this bound, define $T_p = \{(a, b, c) \in \mathbb{Z}^3 : a, b, c \geq 0, a + b + c = p - 1\}$.

Given a probability-distribution π on T_p , define its *marginal* $\mu(\pi)$ to be the probability distribution on $\mathbb{Z}$ given by

$$\mu(\pi)(a) = \sum_{(a,b,c) \in T} \pi(a, b, c).$$

Given a probability distribution π on any finite set S , define the *entropy* of π to be

$$\eta(\pi) = - \sum_{i \in S} \pi(i) \log \pi(i).$$

Let Δ_p be the set of all probability distributions on $\mathbb{Z}$ that arise as the marginal $\mu(\pi)$ for S_3 -symmetric distributions π on T_p . Then if we define $\lambda_p = \max\{\eta(\pi) : \pi \in \Delta_p\}$, Kleinberg, Sawin and Speyer [3] showed that:

Theorem 1. *Let p be a prime, and define λ_p as above. For any n , all sum-free sets are of size at most $e^{\lambda_p n}$.*

For n sufficient large, there are sum-free sets of size at least

$$e^{\lambda_p n - 2\sqrt{2\lambda \log 2\sqrt{n} - O_p(\log n)}}.$$

There is a natural comparison between this and the bound of Ellenberg and Gijswijt [2]. Clearly any distribution π in Δ_p satisfies:

- (1) The expected value of π is $\frac{p-1}{3}$.
- (2) π takes values in $\{0, 1, \dots, p-1\}$.

Let Δ'_p be the larger set of all probability distributions on $\mathbb{Z}$ that have support inside $\{0, 1, \dots, p-1\}$ and have expected value $\frac{p-1}{3}$. Then the problem of finding $\lambda'_p = \max\{\eta(\pi) : \pi \in \Delta'_p\}$ is considerably easier.

Let ρ be the unique positive real number such that the probability distribution

$$\psi_\rho(j) = \frac{\rho^j}{1 + \rho + \dots + \rho^{q-1}} : 0 \leq j \leq q-1$$

has expected value $\frac{p-1}{3}$. Then ψ_ρ is the unique distribution in Δ'_p with maximal entropy.

Ellenberg and Gijswijt showed that

Theorem 2. *Let p be a prime, and let $\lambda'_p = \eta(\psi_\rho)$ be as defined above.*

Then all sets in $\mathbb{Z}_p^n$ with no three-term arithmetic progression are of size at most $Ce^{\lambda'_p n}$.

Kleinberg, Sawin and Speyer [3] conjectured that this distribution ψ_ρ is in Δ_p , and so these two bounds are the same.

The purpose of this note is to close this gap. Define a distribution π in Δ'_p to be *bottom-heavy* if $\pi(0) \geq \pi(1) \geq \dots \geq \pi(p-1)$.

Our proof will not require p to be prime, and will proceed by an induction through all integers. Note that none of the definitions so far depend on p being prime. We will prove the following:

Theorem 3. *For any bottom-heavy distribution π in Δ'_p (for any positive integer $p \geq 1$), π is in Δ_p .*

Since ψ_ρ is clearly bottom-heavy, it follows that ψ_ρ is in Δ_p and hence the Ellenberg-Gijswijt bound and the Kleinberg-Sawin-Speyer bound are the same.

2. METHODOLOGY

Let us say that a *scaled probability distribution* ϕ on the finite set S is a mapping $\phi : S \rightarrow \mathbb{R}^+$ which is not identically 0, and say the *expected value* of scaled probability distribution ϕ is

$$\frac{\sum_{i \in S} i\phi(i)}{\sum_{i \in S} \phi(i)}.$$

We view the scaled probability distribution ϕ as being equal to the ordinary probability distribution

$$\bar{\phi}(i) = \frac{\phi(i)}{\sum_{i \in S} \phi(i)}$$

scaled up by $\sum_{i \in S} \phi(i)$.

Given an element $t = (i, j, k) \in T_p$, denote by η_t the scaled probability distribution on $\{0, 1, \dots, p-1\}$ of the form

$$\eta_t(r) = \mathbb{1}_{r=i} + \mathbb{1}_{r=j} + \mathbb{1}_{r=k}.$$

Say that a scaled probability distribution is *simple* if it of the form η_t .

Lemma 4. *For a scaled probability distribution ϕ , the following are equivalent:*

- (1) $\bar{\phi}$ is the marginal of an S_3 -symmetric distribution on T_p
- (2) ϕ is the positive linear combination of simple distributions.

Proof. If ϕ can be written as $\phi = \sum_{t \in T} \lambda_t \eta_t$, then if we choose an element of T with probability $\frac{\lambda_t}{\sum_t \lambda_t}$ and permute the coordinates uniformly at random, we have an S_3 -symmetric distribution with marginal $\bar{\phi}$.

Similarly if $\bar{\phi}$ is the marginal of some S_3 symmetric distribution ψ on T_p , $\bar{\phi} = \frac{1}{3} \sum_{t \in T_p} \psi_t \eta_t$, so $\phi = \sum_{t \in T_p} \frac{\sum_{i \in S} \phi(i) \psi_t}{3} \eta_t$. $\square$

We extend the definition above of bottom-heavy and of Δ'_p to scaled probability distributions, so Δ'_p is the set of maps $\pi : \{0, 1, \dots, p-1\} \rightarrow \mathbb{R}^+$ that are not identically zero and satisfy $\pi(1) + 2\pi(2) + \dots + (p-1)\pi(p-1) = \frac{p-1}{3}(\pi(0) + \dots + \pi(p-1))$.

If $0 \leq i \leq \frac{2(p-1)}{3} \leq j \leq p-1$ with $i < j$, define the *cornerstone distribution* $\gamma_{i,j,p}$ by

$$\gamma_{i,j,p}(k) = \begin{cases} (j+1)(3j-2(p-1)) + (i+1)(2(p-1)-3i) & \text{if } k \leq i \\ (i+1)(2(p-1)-3i) & \text{if } i < k \leq j \\ 0 & \text{if } j < k. \end{cases}$$

Note that the values $(j+1)(3j-2(p-1))$ and $(i+1)(2(p-1)-3i)$ are of exactly the correct ratio to make the expected value of $\gamma_{i,j,p}$ be $\frac{p-1}{3}$. In Section 3 we will show that any bottom-heavy scaled probability distribution in Δ'_p is the linear combination of cornerstone distributions. Sections 4

to 6 of the paper will be dedicated to showing inductively on p that all cornerstone distributions are the linear combination of simple distributions.

To make the inductive step, the following definitions will be useful. Say that scaled distribution π is *reducible* if $\pi(0) = \pi(p-2) = \pi(p-1) = 0$ and $\pi(1) \geq \pi(2) \geq \dots \geq \pi(p-3)$, and is *flippable* if there exists $i < j$ with $j < \frac{2(p-1)}{3}$ and with $p-1 \leq 2j+i$ such that the support of π is $\{i, i+1, \dots, j\}$ and $\pi(i) \leq \pi(i+1) \leq \dots \leq \pi(j)$.

By exhaustive case analysis, we will prove the following Theorem:

Theorem 5. *For all integers i, j, p with $0 \leq i \leq \frac{2(p-1)}{3} \leq j \leq p-1$, the cornerstone distribution $\gamma_{i,j,p}$ can be expressed as a positive linear combination of simple distributions, reducible distributions and flippable distributions.*

Finally, in Section 7, we will show that this implies Theorem 3

3. CORNERSTONE DISTRIBUTIONS

Our methodology will be to show that cornerstone distributions can be expressed as the linear combination of simpler items. We note here that this is sufficient to prove our main Theorem.

Theorem 6. *Any scaled bottom-heavy probability distribution taking values in $\{0, 1, \dots, p-1\}$ with expected value $\frac{p-1}{3}$ can be expressed as the linear combination of cornerstone distributions.*

Proof. For such probability distribution π , let i be the minimum value in $\{0, 1, \dots, p-1\}$ such that $\pi(i) \neq \pi(i+1)$ and let j be the maximum value.

We prove by induction on $j-i$. Let $j-i = t \geq 0$ and suppose the statement is true for all bottom-heavy probability distributions π' with expected value $\frac{p-1}{3}$ and smaller $j-i$.

We note first that the expected value of π is at least $\frac{i}{2}$:

$$\begin{aligned} \mathbb{E}\pi &= \sum_{t=0}^{p-1} \pi(t)t \\ &\leq \sum_{t=0}^i \pi(0)t + \sum_{t=i+1}^{p-1} \pi(t)\frac{i}{2} \\ &= \pi(0)(i+1)\frac{i}{2} + \frac{i}{2} \sum_{t=i+1}^{p-1} \pi(t) \\ &= \frac{i}{2} \left(\sum_{t=0}^{p-1} \pi(t) \right) = \frac{i}{2}. \end{aligned}$$

Similarly, we note this expected value is at most $\frac{j}{2}$:

$$\mathbb{E}\pi = \sum_{t=0}^{p-1} \pi(t)t$$

$$\begin{aligned}
&= \sum_{t=0}^j \pi(t)t \\
&= \sum_{t < \frac{j}{2}} \pi(t)t + \pi(j-t)t + \sum_{t=\frac{j}{2}} \pi(t)t \\
&\leq \sum_{t < \frac{j}{2}} \pi(t)\frac{j}{2} + \pi(j-t)\frac{j}{2} + \sum_{t=\frac{j}{2}} \pi(t)t \\
&= \sum_{t=0}^j \pi(t)\frac{j}{2} = \frac{j}{2}.
\end{aligned}$$

Thus i and j are in the region for which the cornerstone distribution $\gamma_{i,j,p}$ is well-defined, and in this region it can easily be checked that it is in fact a scaled probability distribution, that it has mean $\frac{p-1}{3}$ and only has $\gamma_{i,j,p}(k) \neq \gamma_{i,j,p}(k+1)$ for $k = i, j$.

Thus for all $0 \leq \alpha < 1$, if we define $\tau_\alpha(k)$ by

$$\tau_\alpha(k) = \frac{\pi(k) - \alpha\gamma_{i,j,p}(k)}{1 - \alpha},$$

then we have $\sum \tau(k) = 1$, $\sum \tau(k)k = \frac{p-1}{3}$, and $\tau_\alpha(k) \geq \tau_\alpha(k+1)$ if $k \neq i, j$.

Choosing the maximal α where $\tau_\alpha(i) \geq \tau_\alpha(i+1)$ and $\tau_\alpha(j) \geq \tau_\alpha(j+1)$ expresses π as the linear combination of α times a cornerstone distribution and $1 - \alpha$ times a bottom-heavy distribution of mean $\frac{p-1}{3}$ with a smaller $j - i$. $\square$

4. A SIMPLE CASE

In the case where $i = \frac{2(p-1)}{3}$, it is particularly easy to express $\gamma_{i,j,p}$ as the linear combination of simple distributions.

Theorem 7. *For $0 \leq i \leq \frac{2(p-1)}{3} \leq j \leq p-1$ with $i < j$, if $i = \frac{2(p-1)}{3}$ or $j = \frac{2(p-1)}{3}$, the cornerstone distribution $\gamma_{i,j,p}$ is in Δ_p .*

Proof. Let $k = \frac{p-1}{3}$. Since k is equal to either $\frac{i}{2}$ or $\frac{j}{2}$, k is a fraction with denominator 2 and a fraction with denominator 3, so k is an integer, $p = 3k + 1$ and $i = 2k$ or $j = 2k$. It is easy to check that for any value of the other parameter i or j , $\gamma_{i,j,p}$ is uniform on $\{0, 1, \dots, 2k\}$.

Three copies of $\{0, 1, \dots, 2k\}$ can easily be split into k triples adding to $3k$: $(i, i+k, 2k-i)$ for $0 \leq i \leq k$ and $(i, i-(k+1), (4k+1)-2i)$ for $k+1 \leq i \leq 2k$. Note that if we select one of these triples uniformly at random, each coordinate is uniform on $\{0, 1, \dots, 2k\}$.

Then $\gamma_{i,j,p}$ is a multiple of the sum of the simple distributions defined by these triples, and hence is in Δ_p . $\square$

5. THE CASE $j < p - 2$

Recall that π is *reducible* if $\pi(0) = \pi(p-2) = \pi(p-1) = 0$ and $\pi(1) \geq \pi(2) \geq \dots \geq \pi(p-3)$.

Theorem 8. *For $0 \leq i < \frac{2(p-1)}{3}$ and $\frac{2(p-1)}{3} \leq j < p-2$, the cornerstone distribution $\gamma_{i,j,p}$ is the linear combination of some simple distributions and a reducible distribution.*

Proof. Given a scaled distribution τ with $\tau(1) \geq \tau(2) \geq \dots \tau(p-1)$ and $\mathbb{E}\tau = \frac{p-1}{3}$, let j be the largest element of the support of τ , let x be equal to $\min(\frac{\tau(0)}{2(j+1)-p}, \frac{\tau(j)}{2})$ and let $\tau' = \tau - x \sum_{t=p-1-j}^j \eta_{0,t,p-1-t}$.

Then it can be seen that we are subtracting $(2(j+1)-p)x$ from $\tau(0)$ and $2x$ from $\tau(t)$ for all t with $p-1-j \leq t \leq j$. As such, τ' is a scaled distribution with $\tau'(1) \geq \tau'(2) \geq \dots \tau'(p-1)$ and $\mathbb{E}\tau' = \frac{qp-1}{3}$.

Let us apply this operator to $\gamma_{i,j,p}$. If $\gamma'_{i,j,p}(0) = 0$, then $\gamma_{i,j,p}$ is the linear combination of simple distributions $\eta_{0,t,p-1-t}$ and a scaled reducible (since $j < p-2$) distribution $\gamma'_{i,j,p}$.

If not, then we have decreased all values in the region $p-1-j \leq t \leq j$ by $\gamma_{i,j,p}(j)$. In particular, if $k = \max(i, p-1-j)$, $\gamma'_{i,j,p}(k) = 0$, so the support of $\gamma'_{i,j,p}$ is contained in $[0, k]$. Further, $\gamma'_{i,j,p}$ is bottom-heavy on $[1, k]$, so the expected value is at most $\frac{k+1}{2}$. Since we know $\gamma'_{i,j,p}$ has mean $\frac{p-1}{3}$, it follows that $k+1 \geq \frac{2(p-1)}{3}$.

Since $p-1-j \leq \frac{p-1}{3}$ and $i \leq \frac{2(p-1)}{3}$, it follows that $k = i$ and $i = \lfloor \frac{2(p-1)}{3} \rfloor$. It follows also that if we apply the above operator again, we will still have $\mathbb{E}\gamma''_{i,j,p} = \frac{p-1}{3}$, so the support of $\gamma''_{i,j,p}$ must still reach as high as i , so $\gamma''_{i,j,p}(0)$ must be 0. Then $\gamma'_{i,j,p}$ is the linear combination of simple distributions and a scaled reducible distribution, and therefore so is $\gamma_{i,j,p}$. $\square$

6. THE CASE $j \geq p-2$

To prove the case $j \geq p-2$, we will first show the existence of some particular distributions in Δ_p . We will use the following Lemma.

Lemma 9. *For integers $m, n \geq 0$, one can pair together random variables X and Y , uniform on $[m]$ and $[n]$ respectively, such that $X+Y$ is uniform on $[m+n]$.*

Similarly, for $m, n \geq 1$, one can pair together random variables X and Y , uniform on $[m]$ and $[n]$ respectively, such that $X+Y$ is uniform on $[1, m+n-1]$.

Proof. For the first part, let $A_1, \dots, A_m, B_1, \dots, B_n$ and C be independent random variables drawn from the uniform distribution $U[0, 1]$. Let $X = |\{i : A_i \leq C\}|$ and $Y = |\{i : B_i \leq C\}|$. Then it is clear that X, Y and $X+Y$ are uniform on $[m]$, $[n]$ and $[m+n]$ respectively.

For the proof of the second part, write $t = m + n$ and define $a_{x,y}$ for $0 \leq x \leq m$ and $0 \leq y \leq n$ by

$$a_{x,y} = \begin{cases} 0 & \text{if } x + y = 0 \text{ or } t \\ \frac{(x+y-1)!(t-x-y-1)!(nx-my)^2}{x!y!(m-x)!(n-y)!} & \text{otherwise.} \end{cases}$$

Clearly the values of $a_{x,y}$ are non-negative. Further, since

$$nx - my \frac{(x+y-1)!}{x!y!} = n \binom{x+y-1}{y} - m \binom{x+y-1}{x}$$

and

$$nx - my \frac{(t-x-y-1)!}{(m-x)!(n-y)!} = m \binom{t-x-y-1}{m-x} - n \binom{t-x-y-1}{n-y},$$

we can rewrite the expression as

$$\begin{aligned} a_{x,y} &= nm \binom{x+y-1}{y} \binom{t-x-y-1}{m-x} - n^2 \binom{x+y-1}{y} \binom{t-x-y-1}{n-y} \\ &\quad - m^2 \binom{x+y-1}{x} \binom{t-x-y-1}{m-x} + mn \binom{x+y-1}{x} \binom{t-x-y-1}{n-y}. \end{aligned}$$

Then, for any fixed x

$$\begin{aligned} \sum_{x+y=r} a_{x,y} &= \sum_{x+y=r} nm \binom{x+y-1}{y} \binom{t-x-y-1}{n-y-1} \\ &\quad - \sum_{x+y=r} n^2 \binom{x+y-1}{y} \binom{t-x-y-1}{n-y} \\ &\quad - \sum_{x+y=r} m^2 \binom{x+y-1}{y-1} \binom{t-x-y-1}{n-y-1} \\ &\quad + \sum_{x+y=r} mn \binom{x+y-1}{y-1} \binom{t-x-y-1}{n-y} \\ &= 2nm \binom{t-1}{n} - n^2 \binom{t-1}{n+1} - m^2 \binom{t-1}{n-1}. \end{aligned}$$

Similarly, for any fixed y

$$\sum_{x=0}^m a_{x,y} = 2nm \binom{t-1}{m} - m^2 \binom{t-1}{m+1} - n^2 \binom{t-1}{m-1}.$$

Finally, for $1 \leq r \leq m+n-1$,

$$\begin{aligned} \sum_{y=0}^n a_{x,y} &= \sum_{y=0}^n nm \binom{r-1}{y} \binom{t-r-1}{n-y-1} \\ &\quad - \sum_{y=0}^n n^2 \binom{r-1}{y} \binom{t-r-1}{n-y} \end{aligned}$$

$$\begin{aligned}
& - \sum_{y=0}^n m^2 \binom{r-1}{y-1} \binom{t-r-1}{n-y-1} \\
& + \sum_{y=0}^n mn \binom{r-1}{y-1} \binom{t-r-1}{n-y} \\
& = 2nm \binom{t-2}{n-1} - n^2 \binom{t-2}{n} - m^2 \binom{t-2}{n-2}.
\end{aligned}$$

Thus if a random variable (x, y) is picked with scaled distribution $a_{x,y}$, x is uniform on $[m]$, y is uniform on $[n]$ and $x+y$ is uniform on $[1, m+n-1]$. $\square$

For integers m, n with $m+n < p$, define scaled distribution $\alpha_{m,n,p}$ by

$$\begin{aligned}
\alpha_{m,n,p} &= (n+1)(m+n+1)\mathbb{1}_{[m]} \\
&+ (m+1)(m+n+1)\mathbb{1}_{[n]} \\
&+ (m+1)(n+1)\mathbb{1}_{[p-1-(m+n), p-1]}.
\end{aligned}$$

Similarly, for integers m, n with $1 \leq m, n$ and $m+n \leq p$, define scaled distribution $\beta_{m,n,p}$ by

$$\begin{aligned}
\beta_{m,n,p} &= (n+1)(m+n-1)\mathbb{1}_{[m]} \\
&+ (m+1)(m+n-1)\mathbb{1}_{[n]} \\
&+ (m+1)(n+1)\mathbb{1}_{[p-(m+n), p-2]}.
\end{aligned}$$

Then the way we will use Lemma 9 is to show that these distributions are in Δ_p .

Corollary 10. *For integers m, n with $m+n < p$, the scaled distribution $\alpha_{m,n,p}$ is in Δ_p and for integers m, n with $1 \leq m, n$ and $m+n \leq p$, the scaled distribution $\beta_{m,n,p}$ is in Δ_p .*

Proof. By the first part of Lemma 9, there is a random distribution (X, Y, Z) on T_p such that X is uniform on $[m]$, Y is uniform on $[n]$ and $Z = (p-1) - X - Y$ is uniform on $[p-1-(m+n), p-1]$.

If we randomly permute the coordinates of this distribution, we get an S3-symmetric distribution on T_p with marginal

$$\frac{\alpha_{m,n,p}}{3(m+1)(n+1)(m+n+1)}.$$

Similarly the second part of Lemma 9 gives rise to an S3-symmetric distribution on T_p with marginal

$$\frac{\beta_{m,n,p}}{3(m+1)(n+1)(m+n-1)}.$$

$\square$

This allows us to simply deal with all remaining cases with a fixed i and sufficiently large p .

Theorem 11. For $0 \leq i \leq \frac{2(p-1)}{3} \leq j \leq p-1$ with $i < j$, if $j \geq p-2$ and $p \geq 2i+2$, $\gamma_{i,j,p}$ is in Δ_p .

Proof. **Case I:** $j = p-1, 2i+2 \leq p$ We claim that

$$\gamma_{i,j,p} = \frac{p-(i+1)}{i+1} \alpha_{i,i,p} + \alpha_{i,p-(2i+2),p}$$

Note that

$$\begin{aligned} \gamma_{i,j,p} &= \gamma_{i,p-1,p} \\ &= (i+1)(2(p-1)-3i) + (p-1)p \mathbb{1}_{[i]} \\ \alpha_{i,i,p} &= 2(i+1)(2i+1) \mathbb{1}_{[i]} + (i+1)^2(1 - \mathbb{1}_{[p-(2i+2)]}) \\ \alpha_{i,p-(2i+2),p} &= (p-(2i+1))(p-(i+1)) \mathbb{1}_{[i]} + (i+1)(p-(i+1)) \mathbb{1}_{[p-(2i+2)]} \\ &\quad + (p-(2i+1))(i+1)(1 - \mathbb{1}_{[i]}). \end{aligned}$$

The equality comes from equating constant terms

$$(i+1)(2(p-1)-3i) = \frac{p-(i+1)}{i+1} (i+1)^2 + (i+1)(p-(2i+1)),$$

$\mathbb{1}_{[i]}$ terms

$$\begin{aligned} (p-1)p &= \frac{p-(i+1)}{i+1} 2(i+1)(2i+1) \\ &\quad + (p-(2i+1))(p-(i+1)) - (p-(2i+1))(i+1), \end{aligned}$$

and $\mathbb{1}_{[p-(2i+2)]}$ terms

$$0 = -\frac{p-(i+1)}{i+1} (i+1)^2 + (i+1)(p-(i+1)).$$

Case II: $j = p-2, 2i+2 \leq p \leq 3i+1$ Similarly

$$\gamma_{i,j,p} = \frac{j-i}{i+1} \beta_{i,i,p} + \beta_{i,p-(2i+1),p}$$

This can be proved in a similar way: each is a linear combination of $1 - \mathbb{1}_{\{p-1\}}$, $\mathbb{1}_{[i]}$, and $\mathbb{1}_{[p-(2i+1)]}$, and the coefficients are equal on both sides. $\square$

We are left to deal with cases with $j \geq p-2$ and $p \leq 2i+1$. Recall that a distribution π in Δ'_p is *flippable* if there exists $i < j$ such that the support of π is $\{i, i+1, \dots, j\}$ and $\pi(i) \leq \pi(i+1) \leq \dots \leq \pi(j)$ with $j < \frac{2(p-1)}{3}$ and with $p-1 \leq 2j+i$.

Theorem 12. For $0 \leq i < \frac{2(p-1)}{3} \leq j \leq p-1$ with $i < j$, if $j \geq p-2$ and $p \leq 2i+1$, $\gamma_{i,j,p}$ is a linear combination of simple distributions and a flippable distribution.

Proof. For now, suppose that $j = p-1$. Let $k = \lfloor \frac{p-2-i}{2} \rfloor$ and $l = \lceil \frac{p-2-i}{2} \rceil$.

Then since $(p-1) - (k+l) = (p-1) - (p-2-i) = i+1 > l$, the scaled distribution $\alpha_{k,l,p}$ takes values

$$\alpha_{k,l,p}(t) = \begin{cases} (k+l+1)(k+l+2) & \text{if } t \leq k \\ (k+1)(k+l+1) & \text{if } k < t \leq l \\ 0 & \text{if } l < t \leq i \\ (k+1)(l+1) & \text{if } i+1 < t \leq p-1. \end{cases}$$

While $\gamma_{i,j,p}$ takes values

$$\gamma_{i,j,p}(t) = \begin{cases} p(p-1) + (i+1)(2(p-1) - 3i) & \text{if } t \leq i \\ (i+1)(2(p-1) - 3i) & \text{if } i+1 < t \leq p-1. \end{cases}$$

Further, we state that $\gamma_{i,j,p}(0)/\alpha_{k,l,p}(0) \geq \gamma_{i,j,p}(p-1)/\alpha_{k,l,p}(p-1)$, which we'll prove later.

It follows that scaled distribution

$$\pi = \gamma_{i,j,p} - \frac{\gamma_{i,j,p}(p-1)}{\alpha_{k,l,p}(p-1)}$$

has $\pi(0) \leq \pi(1) \leq \dots \leq \pi(i)$ and $\pi(i+1) = \dots = \pi(p-1) = 0$. Thus, since $i < 2\frac{p-1}{3}$ and $p \geq 2i+1$, π is flippable.

Similarly, for $j = p-2$, if we let $k = \lfloor \frac{p-1-i}{2} \rfloor$ and $l = \lceil \frac{p-1-i}{2} \rceil$, we can remove a multiple of $\beta_{k,l,p}$ from $\gamma_{i,j,p}$ and get a flippable distribution. $\square$

7. PUTTING IT ALL TOGETHER

We have now covered all cases of Theorem 5

Proof of Theorem 5. Fix integers i, j, p with $0 \leq i \leq \frac{2(p-1)}{3} \leq j \leq p-1$, and we will show that we have showed that $\gamma_{i,j,p}$ can be expressed as a combination of simple distributions, reducible distributions and flippable distributions.

Firstly if $i = \frac{2(p-1)}{3}$ or $j = \frac{2(p-1)}{3}$, we showed in Theorem 7 that $\gamma_{i,j,p}$ was in Δ_p and so a combination of simple distributions.

If $i < \frac{2(p-1)}{3}$ and $j < p-2$, we showed in Theorem 8 that $\gamma_{i,j,p}$ was a combination of simple distributions and a reducible distribution.

If $i < \frac{2(p-1)}{3}$ and $p-2 \leq j$ and $p \geq 2i+2$, we showed in Theorem 11 that $\gamma_{i,j,p}$ was in Δ_p and so a combination of simple distributions.

Finally, if $i < \frac{2(p-1)}{3}$ and $p-2 \leq j$ and $p \leq 2i+1$, we showed in Theorem 12 that $\gamma_{i,j,p}$ was a combination of simple distributions and a flippable distribution.

Clearly this exhausts all cases. $\square$

We now have enough to prove the main Theorem of this note.

Proof of Theorem 3. We prove by induction on p . For $p = 1, 2$ there is only one probability distribution π on $\{0, 1, \dots, p-1\}$ with expected value $\frac{p-1}{3}$.

For $p = 1$ this is $\pi(0) = 1$, which is clearly the marginal of the only S_3 -symmetric distribution $\pi((0, 0, 0)) = 1$ on T .

For $p = 2$ this is $\pi(0) = \frac{2}{3}$, $\pi(1) = \frac{1}{3}$, which is clearly the marginal of the only S_3 -symmetric distribution $\pi(e_1) = \pi(e_2) = \pi(e_3) = \frac{1}{3}$ on T .

Now suppose $p \geq 3$ and the statement is true for all smaller p . Let π be a bottom-heavy distribution on $\{0, 1, \dots, p-1\}$ with mean $\frac{p-1}{3}$.

Then by Theorem 6, π can be expressed as a linear combination of cornerstone distributions.

By Theorem 5, each cornerstone distribution can be expressed as a linear combination of simple distributions, reducible distributions and flippable distributions. Therefore, π can be expressed as a linear combination of simple distributions, reducible distributions.

For any reducible distribution τ , define distribution τ' on $\{0, 1, \dots, p-4\}$ by $\tau'(i) = \tau(i+1)$. Then by the definition of reducibility, τ' is a bottom-heavy distribution on $\{0, 1, \dots, p-4\}$ with mean $\frac{p-4}{3}$. Since the statement is true for $p-3$, τ' is the marginal of an S_3 -symmetric distribution on T_{p-3} . Adding 1 to every coordinate gives an S_3 -symmetric distribution on T_p with marginal τ .

For any flippable distribution τ with parameters $i < j$, define distribution τ' on $\{0, 1, \dots, (3j - (p-1))\}$ by $\tau'(k) = \tau(j-k)$ for $0 \leq k \leq j-i$ and $\tau'(k) = 0$ for $k > j-i$. The requirements on the parameters are $j < \frac{2(p-1)}{3}$, which means that the upper bound on the distribution $(3j - (p-1))$ is less than $p-1$ and $p-1 \geq 2j+i$, which means that $j-i$, the largest element of the support of τ' is less than $3j - (p-1)$. Clearly τ' is a bottom-heavy distribution on $\{0, 1, \dots, (3j - (p-1))\}$ with mean $j - \frac{p-1}{3}$. Since the statement is true for $3j+2-p$, τ' is the marginal of an S_3 -symmetric distribution on T_{3j+2-p} . Subtracting every coordinate from j gives an S_3 -symmetric distribution on T_p with marginal τ .

As noted earlier, simple distributions are clearly the marginal of an S_3 -symmetric distribution on T .

As such, π can be expressed as a linear combination of marginals of S_3 -symmetric distributions on T , and is therefore itself the marginal of an S_3 -symmetric distribution on T . $\square$

Just to state the implication:

Corollary 13. *The Ellenberg-Gijswijt bound [2] and the Kleinberg-Sawin-Speyer bound [3] are the same.*

Proof. The Ellenberg-Gijswijt bound comes from the fact that the distribution ψ_ρ defined by

$$\psi_\rho(j) = \frac{\rho^j}{1 + \rho + \dots + \rho^{p-1}}$$

(where ρ is the unique real number with $\mathbb{E}\psi_\rho = \frac{p-1}{3}$) is the distribution on $\{0, 1, \dots, p-1\}$ with mean $\frac{p-1}{3}$ with the greatest entropy.

Since this is clearly bottom heavy (if $\rho > 1$, $\mathbb{E}\psi_\rho > \frac{p-1}{2}$), by Theorem 3, ψ_ρ is the marginal of an S_3 -symmetric distribution of T and so is in the domain of the Kleinberg-Sawin-Speyer bound, and hence the two bounds are the same. $\square$

REFERENCES

- [1] J. Blasiak, T. Church, H. Cohn, J. A. Grochow, and C. Umans. On cap sets and the group-theoretic approach to matrix multiplication. *ArXiv e-prints*, May 2016.
- [2] J. S. Ellenberg and D. Gijswijt. On large subsets of F_q^n with no three-term arithmetic progression. *ArXiv e-prints*, May 2016.
- [3] R. Kleinberg, W. F. Sawin, and D. E. Speyer. The Growth Rate of Tri-Colored Sum-Free Sets. *ArXiv e-prints*, June 2016.